

University of California at San Diego – Department of Physics – Prof. John McGreevy
Physics 230 Quantum Phases of Matter, Spr 2024
Assignment 4

Due 11pm Thursday, May 2, 2024

1. **A simple avatar of the Lieb-Schulz-Mattis theorem.** Consider the effective theory describing a system living in the continuum that spontaneously forms a solid, say a cubic lattice in d dimensions. Since translation symmetry is spontaneously broken, the degrees of freedom must include a collection of Goldstone bosons θ^I , where $I = 1..d$ runs over the spatial dimensions. $\theta^I(x)$ is the shift of the atom at location x in the I direction relative to its equilibrium position. These fields live on a circle, because if I shift all the atoms by the lattice spacing, I get back the original lattice.

- (a) Convince yourself that the effective action takes the form

$$S_{\text{elastic}}[\theta^I] = \int d^{d+1}x \kappa^{ijKL} \partial_i \theta^K \partial_j \theta^L + \text{terms with more derivatives}, \quad (1)$$

where the coupling constant κ^{ijKL} is the elasticity tensor. With various symmetries imposed, it can be decomposed further into various tensors with names from the 19th century. These tensors describe things like bending moduli – the rigidity of the solid to various kinds of strain.

- (b) Now suppose that the number of atoms is a conserved quantity. That is, consider a situation where there is also a $U(1)$ symmetry. So we can couple the system to a background gauge field A_μ for this $U(1)$ symmetry. We'll assume this $U(1)$ symmetry is not spontaneously broken. What are the leading terms in the (local!) effective action $S_{\text{eff}}[\theta^I, A_\mu]$ that preserve gauge invariance and translation symmetry?
- (c) Consider the case of $d = 1$. In addition to the terms involving dA , one interesting term is

$$S_\nu[\theta, A] \equiv \frac{\nu}{2\pi} \int A \wedge d\theta = \frac{\nu}{2\pi} \int dx dt A_\mu \partial_\nu \theta \epsilon^{\mu\nu}. \quad (2)$$

One point to notice about it is that it is not obviously gauge invariant, because it depends explicitly on A and not just the gauge-invariant object F . Show that e^{iS_ν} is gauge invariant if ν is an integer.

- (d) What does the new term (2) do? Well, the first question we should ask about an effective action for a background gauge field is: what is the resulting charge density:

$$\rho(x) = \frac{\delta S}{\delta A_0(x)} ?$$

Interpret your result.

- (e) What is the analog of (2) in d dimensions? (That is, find a term in d spatial dimensions involving a single power of A and derivatives of the θ^I that can be written without using the metric.) Show that its coefficient ν is quantized to be an integer. What contribution does it make to the density?
- (f) We can identify the goldstone field θ with the phase field describing the displacements of the atoms from their equilibrium positions:

$$u^i(x, t) = \frac{1}{2\pi} a_I^i \theta^I(x, t) - x^i$$

where \vec{a}_I are generators of the lattice Γ . Then the equilibrium configuration is actually $\theta^I(x, t) = K_i^I x^i$ where $K_i^I \left(\frac{a}{2\pi}\right)_I^j = \delta_i^j$, so K_i^I is the matrix whose columns are the reciprocal lattice generators.

- (g) The conclusion you should find by the gauge invariance argument above, under the present assumptions, is that ν , and hence the equilibrium density must be an integer. This is an avatar of the Lieb-Schulz-Mattis-Oshikawa-Hastings (LSMOH) theorem. Now, you may say to yourself, why can't I make a system at some filling which is not an integer? Indeed, I can take 20007 particles and place them in a volume with 20004 unit cells, and the system must have some groundstate. What gives?

2. **Edge modes of CS theory.** Now we return to abelian Chern-Simons theory (for an extra challenge, redo this part in the non-Abelian case). If there is a boundary of spacetime, something must be done to fix up the fact that the action is not invariant under would-be gauge transformations that are nontrivial at the boundary. Consider the case where $\Sigma = \mathbb{R} \times \text{UHP}$ where \mathbb{R} is the time direction, and UHP is the upper half-plane $y > 0$. One way to fix the problem is simply to declare that the would-be gauge transformations which do not vanish at $y = 0$ are not redundancies. This means that they represent physical degrees of freedom.

- (a) First consider the simplest case of U(1) CS theory at level k . Choose $a_0 = 0$ gauge, and plug the solution of the bulk equations of motion $a = \tilde{d}\phi$ (where $\phi(x, y \rightarrow 0) \equiv \phi(x)$ is a scalar field, and \tilde{d} is the exterior derivative on the

spatial manifold) into the Chern-Simons action to find the resulting action for ϕ .

- (b) We can also add local terms at the boundary to the action. Consider adding $\Delta S = g \int_{\partial\Sigma} a_x^2$ (for some coupling constant g). Find the equations of motion for ϕ .

Interpretation: the Chern-Simons theory on a space with boundary necessarily produces a chiral edge mode.

- (c) If you feel like it, redo the previous parts for the general K -matrix theory.