University of California at San Diego – Department of Physics – Prof. John McGreevy Physics 211C (239) Phases of Quantum Matter, Spring 2021

Assignment 6

Due 12:30pm Wednesday, May 19, 2021

Thanks for following the submission guidelines on hw01. Please ask me by email if you have any trouble.

## 1. Boson Integer Quantum Hall State from Partons.

Consider a system made from two species of bosons,  $b_{\uparrow}, b_{\downarrow}$ . They could be two layers. We'll assume that only the total boson number, acting by  $(b_{\uparrow}, b_{\downarrow}) \rightarrow e^{i\alpha}(b_{\uparrow}, b_{\downarrow})$  is conserved, and couple to a background field  $\mathcal{A}$  for that symmetry.

(a) Consider the parton ansatz:

$$b_{\uparrow} = f_0 f_{\uparrow}, \quad b_{\downarrow} = f_0 f_{\downarrow} f_1 f_2$$

where all the fs are fermionic partons. There are three U(1) gauge fields that glue these partons back together, and the charge assignments are as follows:

	$a_1$	$a_2$	$a_3$	$\mathcal{A}$	Chern $\#$ in	Chern $\#$ in	Chern $\#$ in
					Phase 1	Phase 2	Phase 3
$f_{\uparrow}$	1	0	0	1	1	1	1
$f_{\downarrow}$	1	1	0	1	1	1	1
$f_0$	-1	0	0	0	-1	-1	-1
$f_1$	0	-1	1	0	-1	-1	-1
$f_2$	0	0	-1	0	-1	0	1

Also in the table are the Chern numbers of the bands filled by each of the partons in three distinct phases. (Only the Chern number of  $f_2$  changes.) Identify the three phases, and describe the critical theories separating them. Hint: I recommend describing the parton currents in terms of dynamical gauge fields  $j^{(\alpha)}_{\mu} = \frac{1}{2\pi} \epsilon_{\mu\nu\rho} \partial_{\nu} b^{(\alpha)}_{\rho}$ , where  $\alpha = \uparrow, \downarrow, 0, 1, 2$ .

(b) For this part of the problem, let's retreat to the continuum. Consider the simpler parton ansatz:

$$b_{\uparrow} = f_0 f_{\uparrow}, \quad b_{\downarrow} = f_0 f_{\downarrow}$$

where all the fs are fermionic partons. Choose the  $U(1)_{\mathcal{A}}$  to be charges  $q_0 = 2, q_{\uparrow} = -1, q_{\downarrow} = -1$ .

Consider an equal number N of  $b_{\uparrow}$  and  $b_{\downarrow}$  particles, so that the total filling fraction is  $\nu = 2$ . How many  $f_0$  particles are there, and how many  $f_{\downarrow}, f_{\uparrow}$  particles are there?

Write a candidate groundstate wavefunction  $\Psi(r_i^{\uparrow}, r_i^{\downarrow})$  for the bosons.

- (c) Bonus question: why does the simpler ansatz of the previous part produce a wavefunction in the same phase as one of the phases of the first part?
- (d) Actually, here is a simpler description of the same phase diagram, closer to what I said in lecture. Consider a single species of boson, with the simple parton ansatz with  $b = d_1d_2$  in terms of two fermions. Let  $d_1$  and  $d_2$  fill Chern bands with total Chern number  $c_1$  and  $c_2$ . Fix  $c_1 = -1$ . Consider what happens when  $c_2 = 2$ .

Describe the effective field theory of  $d_2$  filling two bands with chern number 1 by introducing two gauge fields each with CS term  $\frac{1}{4\pi}b_adb_a$ .