

Physics 215B QFT Winter 2019 Assignment 3

Due 12:30pm Wednesday, January 30, 2019

1. Brain-warmers.

Consider the field theory with action

$$S[\phi] = \int d^{d+1}x \left(\frac{1}{2}(\partial_\mu\phi\partial^\mu\phi - m^2\phi^2) - \frac{g}{3!}\phi^3 \right).$$

- State the Feynman rules in position space for this theory.
- Draw the diagram which corrects the position-space two-point function of ϕ at order g^2 .
- Find the symmetry factor for this diagram.
- Check by explicit perturbation expansion that the symmetry factor is correct.

2. Propagator corrections in a solvable field theory.

Consider a theory of a scalar field in D dimensions with action

$$S = S_0 + S_1$$

where

$$S_0 = \int d^Dx \frac{1}{2} (\partial_\mu\phi\partial^\mu\phi - m_0^2\phi^2)$$

and

$$S_1 = - \int d^Dx \frac{1}{2} \delta m^2 \phi^2 .$$

We have artificially decomposed the mass term into two parts. We will do perturbation theory in small δm^2 , treating S_1 as an ‘interaction’ term. We wish to show that the organization of perturbation theory that we’ve seen lecture will correctly reassemble the mass term.

- Write down all the Feynman rules for this perturbation theory.
- Determine the 1PI two-point function in this model, defined by

$$i\Sigma \equiv \sum (\text{all 1PI diagrams with two nubbins}).$$

(c) Show that the (geometric) summation of the propagator corrections correctly produces the propagator that you would have used had we not split up $m_0^2 + \delta m^2$.

3. **Reminder.** If you didn't do the bonus problem on the last homework with the Catalan numbers, try it now.

4. **Particle creation by an external source.**

Consider the Hamiltonian

$$H = H_0 + \int d^3x (-j(t, \vec{x})\phi(x))$$

where H_0 is the free Klein-Gordon Hamiltonian, ϕ is the Klein-Gordon field, and j is a c-number scalar function. Here we will outline a calculation of particle-production probabilities (as a functional of j) using operator methods. Alternatively, if you prefer, you can do this whole problem using path integral methods. Even better, do both.

(a) Show that the probability that the source creates *no* particles is given by

$$P(0) = |\langle 0 | \mathcal{T} \{ e^{+i \int d^4x j(x)\phi_I(x)} \} | 0 \rangle|^2.$$

(b) Evaluate the term in $P(0)$ of order j^2 , and show that $P(0) = 1 - \lambda + \mathcal{O}(j^4)$ where

$$\lambda = \int \frac{d^3p}{2E_p} |\tilde{j}(p)|^2.$$

We will show below that $\lambda = \langle N \rangle$ is the mean number of particles created by the source.

(c) Represent the term computed in part 4b as a Feynman diagram. Now represent the whole perturbation series for $P(0)$ in terms of Feynman diagrams. Show that this series exponentiates, so that it can be summed exactly $P(0) = e^{-\lambda}$.

(d) Compute the probability that the source creates one particle of momentum k . Perform this computation first to $\mathcal{O}(j)$ and then to all orders, using the trick of the previous part to sum the series.

(e) Show that the probability of producing n particles is given by the Poisson distribution,

$$P(n) = \frac{1}{n!} \lambda^n e^{-\lambda}.$$

(f) Prove the following facts about the Poisson distribution:

$$\sum_{n=0}^{\infty} P(n) = 1, \quad \langle N \rangle \equiv \sum_{n=0}^{\infty} nP(n) = \lambda,$$

that is, $P(n)$ is a probability distribution, and $\langle N \rangle = \lambda$ as predicted. Compute the fluctuations in the number of particles produced $\langle (N - \langle N \rangle)^2 \rangle$.

5. **A background field.** [This is a bonus problem.]

Consider the following action for a real scalar field Φ :

$$S[\Phi] = \int d^{d+1}x \frac{1}{2} (\partial_\mu \Phi \partial^\mu \Phi - m^2 \Phi^2 - g\phi(x)\Phi^2).$$

The last term here is a cubic coupling between ϕ and Φ . But here we will treat $\phi(x)$ as a fixed background field (analogous to $j(x)$ on previous problems) which acts as a spacetime-dependent mass for the dynamical field Φ .

(a) Show that the two-point Green's function, $G(x, y) \equiv \langle \Omega | \mathcal{T} \Phi(x) \Phi(y) | \Omega \rangle$, satisfies the Schwinger-Dyson equation

$$\delta^{d+1}(x - y) = \mathbf{i} (\partial^2 + m^2 + g\phi(x)) G(x, y). \quad (1)$$

(b) We would like to solve this differential equation. As a warmup, consider the case $g = 0$. Here is a trick: add a fictitious additional time direction T

$$(\partial_T + \mathbf{i} (\partial^2 + m^2)) G(x, y, T) = \delta^{d+1}(x - y) \delta(T) \quad (2)$$

This is just a diffusion equation (in $d+2$ dimensions and with a funny factor of \mathbf{i} !). Show that given a solution to (2), you can find the solution of (1) with $g = 0$ by

$$G(x, y) = \int_0^\infty dT G(x, y, T). \quad (3)$$

(c) Show that the solution to the diffusion equation (2) for an infinitesimal time step T is

$$G(x, y, T) = \frac{1}{(2\pi T)^\alpha} e^{a \frac{(x-y)^2}{2T} + b \frac{m^2}{2} T}. \quad (4)$$

Find α, a, b . Use this to construct the path integral representation

$$G(x, y, T) = \int_{x(0)=x}^{x(T)=y} [Dx] e^{-\mathbf{i} \int_0^T d\tau (\dot{x}^\mu \dot{x}_\mu + m^2)}.$$

- (d) For the case of constant m^2 , the infinitesimal solution (4) actually works for finite T . Show by differentiation that plugging (4) into (3) gives an integral representation of the free Klein-Gordon propagator.
- (e) Now let $g \neq 0$ and suppose that ϕ is slowly varying. Generalize the path integral representation to include the dependence on ϕ .
- (f) Consider a non-relativistic situation, where the spacetime points x and y are separated by a timelike distance large compared to $1/m$. Justify and use stationary-phase methods to show that the dominant contribution to the path integral is a straight-line trajectory between the two points x and y . Evaluate the resulting amplitude as a functional of $\phi(x)$.

This calculation shows that the heavy particle made by the field Φ can be treated as a source for ϕ propagating on a fixed path in spacetime.

- (g) Redo the problem for a *charged* scalar field, Φ in the background of a vector potential A_μ , with

$$S[\Phi] = \int d^{d+1}x \frac{1}{2} (D_\mu \Phi^* D^\mu \Phi - m^2 \Phi^2), \quad D_\mu \Phi \equiv \partial_\mu \Phi - i A_\mu \Phi.$$

It will help to recall that the action of a classical charged particle is $\int d\tau (\dot{x}^2 + \dot{x}^\mu A_\mu(x))$.

6. **Classical Maxwell theory.** Classical electromagnetism follows from the action

$$S[A] = \int d^4x \left(-\frac{1}{4} F_{\mu\nu} F^{\mu\nu} + j^\mu A_\mu \right), \quad \text{where } F_{\mu\nu} = \partial_\mu A_\nu - \partial_\nu A_\mu.$$

- (a) Derive Maxwell's equations as the Euler-Lagrange equations of this action, treating the components $A_\mu(x)$ as the dynamical variables

$$0 = \frac{\delta S[A]}{\delta A_\mu(x)}.$$

Write the equations in standard form by identifying $E^i = -F^{0i}$ and $\epsilon^{ijk} B^k = -F^{ij}$.

- (b) Construct the energy-momentum tensor for this theory. Note that the usual procedure

$$T_\nu^\mu = \frac{\partial \mathcal{L}}{\partial(\partial_\mu \phi)} \partial_\nu \phi - \mathcal{L} \delta_\nu^\mu$$

does not result in a symmetric tensor. To remedy that, we can add to $T^{\mu\nu}$ a term of the form $\partial_\lambda K^{\lambda\mu\nu}$, where $K^{\lambda\mu\nu}$ is antisymmetric in its first two indices. Such an object is automatically divergenceless, so

$$\widehat{T}^{\mu\nu} \equiv T^{\mu\nu} + \partial_\lambda K^{\lambda\mu\nu}$$

is an equally good energy-momentum tensor with the same globally conserved energy and momentum. Show that this construction, with

$$K^{\lambda\mu\nu} = F^{\mu\lambda} A^\nu,$$

leads to an energy-momentum tensor \hat{T} that is symmetric and yields the standard formulae for the electromagnetic energy and momentum densities:

$$\mathcal{E} = \frac{1}{2} (E^2 + B^2), \quad \vec{S} = \vec{E} \times \vec{B}.$$

- (c) [Bonus question] What modification of the action will produce the improved stress tensor as the Noether current for translation symmetry?